



Scopus® doi

Journal of Vibration Engineering

ISSN:1004-4523

Registered



SCOPUS



GOOGLE SCHOLAR



DIGITAL OBJECT
IDENTIFIER (DOI)



IMPACT FACTOR 6.1



Our Website
www.jove.science

SquareSumDegreeDividedByDiameterEnergyofGraphs

KavitaPermi

Department of
Mathematics Presidency Uni
versity Bengaluru-
560059, India.

e-mail: kavitafermi@presidencyuniversity.in

ManasaHS

Department of
Mathematics Presidency Uni
versity Bengaluru-
560059, India.

e-mail: manasaangeetha09@gmail.com

PradeepKumarR

Department of Mathematics
GSSS Institute of Engineering and Technology Mys
suru-570016, India.

e-mail: pradeep.r.mysore@gmail.com

Abstract

The square sum degree divided by diameter matrix $\frac{SSD}{diam}(G)$ of a graph G is a square matrix whose (i, j) th entry is $\frac{d^2 + d_j^2}{diam}$ whenever $i = j$ and otherwise zero. where d_i, d_j is the degree of i^{th} and j^{th} vertex of G . In this paper, we define square sum degree divided by diameter energy $E_{\frac{SSD}{diam}}(G)$ as sum of absolute eigenvalues of $\frac{SSD}{diam}(G)$. Also obtained some bounds on $\frac{SSD}{diam}$ eigenvalues and energy.

Keywords: Square sum degree, Diameter, Eigenvalues, Spectrum and Energy

AMS2010 subject classification: 05C50

1 Introduction

The basic idea of graph theory were born in 1736 with Euler's paper in which he solved the Königsberg bridge problem. In the last decades graph theory has established itself as a worthwhile mathematical discipline and there are many applications of graph theory to a wide variety of subjects which include operation research, Physics, Chemistry, Economics, Genetics, Sociology,

Engineering etc. We can associate several matrices with a graph to record information about

²Correspondingauthor:kavitafermi@presidencyuniversity.in

vertices and how they are interconnected. That is, we can give an algebraic structure to every graph. Many interesting results can be proved about graphs using matrices and other algebraic properties. The main use of algebraic structure is that we can translate properties of graphs into algebraic properties and then use the results and methods of algebra to deduce theorems about graphs. We mainly concentrate on energy of graphs which was introduced by I. Gutman in 1978 [5], which is having direct connection with the total π -

electron energy of a molecule in the quantum chemistry as calculated with the Hückel molecular orbital method. Recently several results on energy related with matrices dealing with degree of vertices and distance between vertices have been studied such as distance energy [7, 9], degree sum energy [6], degree exponent energy [11, 10], degree exponent sum energy [8, 3], degree square sum energy [2, 1, 4] etc. In continuation with this, in order to upgrade, we now introduce the concept of degree square sum distance square energy of a connected graph. The purpose of this paper is to compute the square sum degree divided by diameter matrix denoted by $SSDD(G)$.

2 Square sum degree divided by diameter matrix and its energy

Let G be a connected graph of order n with vertex set $V(G) = (v_1, v_2, \dots, v_n)$. We denote by $d(v_i)$ as the degree of a vertex v_i which is the number of edges incident on it and the distance between two vertices v_i and v_j as d_{ij} , the length of the shortest path joining them. Motivated from previous research, we now define the degree square sum degree divided by diameter matrix of a connected graph G as,

$$b_{ij} = \begin{cases} \frac{d_i^2 + d_j^2}{diam(G)} & \text{if there is a path between } v_i \text{ and } v_j \\ 0 & \text{otherwise} \end{cases}$$

This square sum degree divided by diameter matrix is a symmetric matrix with eigenvalues $\psi_1 \geq \psi_2 \geq \psi_3 \geq \dots \geq \psi_p$.

The characteristic polynomial of $\frac{SSD(G)}{diam}$ is given by $\det|\psi I - \frac{SSD(G)}{diam}|$. The square sum degree divided by diameter energy of the graph G is defined as the sum of absolute values of $\psi_i, i=1, 2, \dots, p$.

$$E \frac{SD}{diam}(G) = \sum_{i=1}^p |\psi_i|.$$

3 Properties of Square sum degree divided by diameter energy

Theorem 3.1. If eigenvalues of $\frac{SSD}{diam}(G)$ are $\psi^+ > \psi^+_{-1} > \psi^+_{-2} > \dots > \psi^+$, then

$$\begin{aligned} 1. \sum_{i=1}^n \psi_i &= 0 \text{ and} \\ 2. \sum_{i=1}^n \psi_i^2 &= 2 \frac{\sum_{i=1}^n d^2 + d_i^{22}}{diam(G)} = 2\Phi \end{aligned}$$

$$\text{where } \Phi = \frac{\sum_{i=1}^n d^2 + d_i^{22}}{diam(G)}.$$

Proof. (1) Since the diagonal entries are zero, the sum of leading diagonal entries of $\frac{SSD}{diam}(G)$ is zero.

$$\sum_{i=1}^n \psi_i = 0.$$

(2) The sum of squares of latent roots of $\frac{SSD}{diam}(G)$ is the sum of latent roots of $[\frac{SSD}{diam}(G)]^2$.

$$\begin{aligned} \sum_{i=1}^n \psi_i^2 &= \sum_{i=1}^n \sum_{j=1}^n u_{ij} u_{ji} \\ &= 0 + 2 \sum_{i < j} (u_{ij})^2 \\ &= 2 \frac{\sum_{i=1}^n d^2 + d_i^{22}}{diam(G)} \end{aligned}$$

$$= 2\Phi.$$

$$diam(G)$$

□

Theorem 3.2. If c_0, c_1 and c_2 are the first three coefficients of characteristic polynomial of $\frac{SSD}{diam}(G)$ matrix, then

$$1. c_0 = 1,$$

$$2. c_1 = 0 \text{ and } 3.$$

$$c_2 = -\Phi.$$

Proof. (i) By definition, $\Gamma(\psi, x) = \det[\psi I - \Phi]$. Therefore $c_0 = 1$.

(ii) $c_1 = (-1)^1 \times \text{trace}(\Gamma) = -1 \times 0 = 0$.

4

$$\begin{aligned}
 \text{(iii) By definition } c_2 &= \sum_{1 \leq i < j \leq p} \begin{vmatrix} u_{ii} & u_{ij} \\ u_{ji} & u_{jj} \end{vmatrix} = \sum_{1 \leq i < j \leq p} (a_{ii}u_{jj} - u_{ij}u_{ji}) \\
 &= \sum_{1 \leq i < j \leq p} u_{ii}u_{jj} - \sum_{1 \leq i < j \leq p} a_{ij}^2 = 0 - \Phi = -\Phi.
 \end{aligned}$$

□

We have the following bounds for $\frac{SSD}{diam}(G)$ using McClelland's inequalities.

Theorem 3.3. Let G be a graph with p vertices, then the upper bound for $\frac{SSD}{diam}(G)$ is

$$E \frac{SSD}{diam}(G) \leq \sqrt{2p}\Phi.$$

Proof. Let $\psi_1 \geq \psi_2 \geq \dots \geq \psi_p$ be the eigenvalues of $\frac{SD}{diam}(G)$, then by Using Cauchy-Schwarz inequality we have,

$$\left(\sum_{i=1}^p u_i v_i \right)^2 \leq \left(\sum_{i=1}^p u_i^2 \right) \left(\sum_{i=1}^p v_i^2 \right).$$

Choose $u_i = 1, v_i = \psi_i$ and by Theorem 3.1

$$\begin{aligned}
 \left(\sum_{i=1}^p \psi_i \right)^2 &\leq \sum_{i=1}^p 1^2 \sum_{i=1}^p \psi_i^2 = p \sum_{i=1}^p \psi_i^2 \\
 \frac{SSD}{diam}(G) &\leq \sqrt{2p}\Phi.
 \end{aligned}$$

Hence

$$E \frac{SSD}{diam}(G) \leq \sqrt{2p}\Phi.$$

□

We present the following lower bounds for $E \frac{SSD}{diam}(G)$.

Theorem 3.4. Let G be a graph with p vertices. If $\tau = \det \frac{SSD}{diam}(G)$ of G ,

then the lower bound is

$$E \frac{SSD}{diam}(G) \geq \sqrt{\frac{2\Phi + p(p-1)\tau^2}{p}}$$

Proof. By definition we have,

$$E \frac{SSD}{diam}(G)^2 = \sum_{i=1}^p \psi_i^2 = \sum_{i=1}^p \psi_i \sum_{j=1}^p \psi_j = \sum_{i=1}^p \psi_i^2 + \sum_{i \neq j} \psi_i \psi_j$$

From the inequality of arithmetic and geometric means

$$\begin{aligned} \frac{1}{p(p-1)} \sum_{i \neq j} \psi_i \psi_j &\geq \sqrt[p(p-1)]{\prod_{i \neq j} \psi_i \psi_j} \\ \text{Therefore } \frac{SSD}{diam}(G)^2 &\geq \sum_{i=1}^p \psi_i^2 + p(p-1) \sqrt[p(p-1)]{\prod_{i=1}^p \psi_i^2} \\ &\geq \sum_{i=1}^p \psi_i^2 + p(p-1) \left(\prod_{i=1}^p \psi_i \right)^{\frac{2}{p-1}} \\ &= \sum_{i=1}^p \psi_i^2 + p(p-1) \left(\prod_{i=1}^p \psi_i \right)^{\frac{2}{p-1}} \\ &= 2\Phi + p(p-1) \tau^{\frac{2}{p-1}} \end{aligned}$$

Hence

$$E \frac{SSD}{diam}(G) \geq \sqrt[p(p-1)]{2\Phi + p(p-1) \tau^{\frac{2}{p-1}}}$$

□

Theorem 3.5. Let r_i and s_i , $1 \leq i \leq p$ be positive real numbers with $M_1 = \max_{1 \leq i \leq p} (r_i)$, $M_2 = \max_{1 \leq i \leq p} (s_i)$, $m_1 = \min_{1 \leq i \leq p} (r_i)$, $m_2 = \min_{1 \leq i \leq p} (s_i)$ then by theorem 9 of [?]

$$\sum_{i=1}^p r_i^2 \sum_{i=1}^p s_i^2 \leq \frac{1}{4} \left(\frac{M_1^2 + M_2^2}{m_1 m_2} \right) \sum_{i=1}^p r_i s_i$$

Theorem 3.6. For a graph G with p vertices let

$|\psi_1|$ and $|\psi_p|$ are the maximum and minimum eigenvalues among all $|\psi_i|$ of $SSD(G)$ respectively, then we have

$$E \frac{SSD}{diam}(G) \geq \sqrt{\frac{8p\Phi |\psi_1| |\psi_p|}{|\psi_1| + |\psi_p|}}$$

6

Proof. Consider a graph G with p vertices let $|\psi_1|$ and $|\psi_p|$ are the maximum and minimum eigenvalues among all $|\psi_i|$ of $^{SSD}_{diam}(G)$ respectively.
From theorem 3.5,

$$\sum_{i=1}^p r_i^2 \sum_{i=1}^p s_i^2 \leq \frac{1}{4} \frac{M_1 M_2}{m_1 m_2} \frac{r_1}{M_1 M_2} \sum_{i=1}^p r_i s_i.$$

Let $r_i = 1, s_i = |\zeta_i|, M_1 M_2 = |\psi_1|, m_1 m_2 = |\psi_p|$ then

$$\sum_{i=1}^p 1^2 \sum_{i=1}^p \psi_i^2 \leq \frac{1}{4} \frac{|\psi_1|}{|\psi_p|} \sum_{i=1}^p 1 |\psi_i|$$

From theorem 3.1

$$p^2 \Phi \leq \frac{1}{4} \frac{(|\psi_1| + |\psi_p|)^2}{|\psi_1| |\psi_p|} E \frac{SSD}{diam}(G)^2,$$

$$E \frac{SSD}{diam}(G)^2 \geq \frac{8p\Phi |\psi_1| |\psi_p|}{(|\psi_1| + |\psi_p|)^2_p}$$

$$E \frac{SSD}{diam}(G) \geq \sqrt{\frac{8p\Phi |\psi_1| |\psi_p|}{|\psi_1| + |\psi_p|}}.$$

□

Theorem 3.7. Let r_i and $s_i, 1 \leq i \leq n$ be non negative real numbers with

$M_1 = \max_{1 \leq i \leq n}(r_i), M_2 = \max_{1 \leq i \leq n}(s_i), m_1 = \min_{1 \leq i \leq n}(r_i), m_2 = \min_{1 \leq i \leq n}(s_i)$ then by theorem 3.1 of

$$\sum_{i=1}^n r_i^2 \sum_{i=1}^n s_i^2 - \sum_{i=1}^n r_i s_i \leq \frac{n^2}{4} (M_1 M_2 - m_1 m_2)^2.$$

Theorem 3.8. For a graph G with p vertices, we have

$$E \frac{SD}{diam}(\Gamma) \geq \frac{r}{2p\Phi - \frac{p^2}{4}(|\psi_1| - |\psi_p|)^2}$$

Proof. Consider a graph G with p vertices let $|\psi_1|$ and $|\psi_p|$ are the maximum and minimum eigenvalues among all $|\psi_i|$ of $^{SSD}_{diam}(G)$ respectively.
From theorem 3.7,

$$\sum_{i=1}^p r_i^2 \sum_{i=1}^p s_i^2 - \sum_{i=1}^p r_i s_i \leq \frac{p^2}{4} (M_1 M_2 - m_1 m_2)^2.$$

Let $r_i=1, s_i=|\psi_i|, M_1M_2=|\psi_i|, m_1m_2=|\psi_p|$, then

$$\sum_{i=1}^p \frac{1}{s_i^2} \psi_i^2 - \sum_{i=1}^p \frac{1}{|\psi_i|} \leq \frac{p^2}{4} (|\psi_1| - |\psi_p|)^2.$$

From theorem 3.1

$$2p\Phi - E \frac{SSD}{diam}(G) \leq \frac{p^2}{4} (|\psi_1| - |\psi_p|)^2,$$

$$E \frac{SSD}{diam}(G) \geq \frac{2p\Phi - \frac{p^2}{4} (|\psi_1| - |\psi_p|)^2}{2}$$

□

Theorem 3.9. Let r_i and $s_i, 1 \leq i \leq p$ be positive real numbers, then by [13]

$$\left| \sum_{i=1}^p r_i s_i - \sum_{i=1}^p r_i \sum_{i=1}^p s_i \right| \leq \mu(p)(A - a)(B - b)$$

where a, b, A and B are real constants, that for each $i, 1 \leq i \leq p, a \leq r_i \leq A$

and $b \leq s_i \leq B$. Further, $\mu(p) = p \left[\frac{1}{2} - \frac{1}{2} \left(\frac{1}{p} \right) \right]$

Theorem 3.10. For a graph G with p vertices, we have

$$E \frac{SSD}{diam}(G) \geq \frac{2p\Phi - \mu(p)(|\psi_1| - |\psi_p|)^2}{2}$$

Proof. Consider a graph G with p vertices let $|\psi_1|$ and $|\psi_p|$ are the maximum and minimum eigenvalues among all $|\psi_i|$ of $SSD(G)$ respectively.

From theorem 3.9,

$$\left| \sum_{i=1}^p r_i s_i - \sum_{i=1}^p r_i \sum_{i=1}^p s_i \right| \leq \mu(p)(A - a)(B - b).$$

Let $r_i=s_i=|\psi_i|, A=B=|\psi_1|, a=b=|\psi_p|$ then

$$p \sum_{i=1}^p |\psi_i|^2 - \sum_{i=1}^p |\psi_i| \leq \mu(p)(|\psi_1| - |\psi_p|)(|\psi_1| - |\psi_p|).$$

From theorem 3.1

$$2p\Phi - E \frac{SSD}{diam}(G) \leq \mu(p)(|\psi_1| - |\psi_p|)^2,$$

$$E \frac{SSD}{diam}(G) \geq \frac{2p\Phi - \mu(p)(|\psi_1| - |\psi_p|)^2}{2}$$

□

4 Squaresumdegreebydiametermatrixanditsenergyforstandardgraphs

Theorem4.1. Let K_p be a complete graph with p vertices, then

$$E \frac{SSD}{diam}(K_p) = 2p^3 + 12p^2 - 82p + 116.$$

Proof. The complete graph K_p with p -vertices has their squaresumdegreebydiameter matrix as follows

$$SSD_{diam}(K_p) = \begin{bmatrix} 0 & 2(p-1)^2 & 2(p-1)^2 & \dots & 2(p-1)^2 \\ 2(p-1)^2 & 0 & 2(p-1)^2 & \dots & 2(p-1)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 2(p-1)^2 & 2(p-1)^2 & 0 & \dots & 2(p-1)^2 \\ \vdots & \vdots & \vdots & \vdots & 0 \end{bmatrix}.$$

Its characteristic polynomial is,

$$[\psi - (18p^2 - 88p + 118)][\psi - (-(2p^2 - 4p + 2))]^{p-1} = 0.$$

Spectra $\frac{SSD}{diam}(K_p) = \begin{matrix} (18p^2 - 88p + 118) & -(2p^2 - 4p + 2) \\ 1 & p-1 \end{matrix}.$

Therefore

$$E \frac{SSD}{diam}(K_p) = |(18p^2 - 88p + 118)|1| + |-(2p^2 - 4p + 2)|(p-1) \\ = 2p^3 + 12p^2 - 82p + 116.$$

□

Theorem4.2. Let S_p^0 , $p \geq 3$ be a crown graph with $2p$ vertices, then

$$E \frac{SSD}{diam}(S_p^0) = \frac{4p^3 + 28p^2 - 172p + 236}{3}.$$

Proof. The crown graph S_p^0 with p -vertices has its squaresumdegreebydiameter matrix as follows

$$SSD_{diam}(S_p^0) = \begin{bmatrix} 0 & \frac{2(p-1)^2}{3} & \frac{2(p-1)^2}{3} & \dots & \frac{2(p-1)^2}{3} \\ \frac{2(p-1)^2}{3} & 0 & \frac{2(p-1)^2}{3} & \dots & \frac{2(p-1)^2}{3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{2(p-1)^2}{3} & \frac{2(p-1)^2}{3} & 0 & \dots & \frac{2(p-1)^2}{3} \\ \vdots & \vdots & \vdots & \vdots & 0 \end{bmatrix}.$$

h Its characteristic polynomial is

$$\psi - \frac{38p^2 - 180p + 238}{3} \quad \psi - \frac{-(2p^2 - 4p + 2)}{3} = 0$$

$$Spectra \frac{SSD}{diam} (S^0_p) = \frac{\frac{38p^2-180p+238}{3}}{1} \frac{\frac{-(2p^2-4p+2)}{3}}{2p-1}.$$

Therefore

$$E_{diam}(S_p) = \frac{38p^2 - 180p + 238}{3} \cdot (1) + \frac{-(2p^2 - 4p + 2)}{3} \cdot (2p - 1)$$

$$= \frac{4p^3 + 28p^2 - 172p + 236}{3}$$

☐

Theorem 4.3. *Let $K_{p \times 2}$ be a cocktail party graph with $2p$ vertices, then*

$$E \frac{SSD}{diam}(K_{p \times 2}) = 8p^3 + 56p^2 - 344p + 472.$$

Proof. The cocktail party graph $K_{p \times 2}$ with $2p$ -vertices has its squaresum degree by diameter matrix as follows

$$SSD_{diam(K)}^{p \times 2} = \begin{pmatrix} 0 & 4(p-1)^2 & 4(p-1)^2 & \dots & 4(p-1)^2 \\ 4(p-1)^2 & 0 & 4(p-1)^2 & \dots & 4(p-1)^2 \\ 4(p-1)^2 & 4(p-1)^2 & 0 & \dots & 4(p-1)^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 4(p-1)^2 & 4(p-1)^2 & \dots & 4(p-1)^2 & 0 \end{pmatrix}.$$

Its characteristic polynomial is

$$[\psi - (76p^2 - 360p + 476)][\psi - (4p^2 - 8p + 4)]^{2p-1} = 0$$

$$Spectra \frac{SSD}{diag}(K_{p \times 2}) = \frac{(76p^2 - 360p + 476)(4p^2 - 8p + 4)1}{2p - 1}.$$

Therefore

$$E \frac{SSD}{diag} (K_p \times 2) = \frac{(76p^2 - 360p + 476) \cdot 1 + 4p^2 - 8p + 4 \cdot (2p - 1)}{1}$$

10

$$=8p^3+56p^2-344p+472.$$

□

Theorem 4.4. Let $K_{p,p}$ be a double star graph with p vertices, then

$$E \frac{SSD}{diam}(K_{p,p}) = \frac{1}{48}(272p^2 - 632p + 756).$$

Proof. The double star graph $K_{p,p}$ with p -vertices has its square sum degree by diameter matrix as follows

$$\begin{array}{cccccccc} \begin{array}{c} \square \\ 0 \end{array} & \begin{array}{c} \frac{p^2+1}{3} \\ \square \end{array} & \begin{array}{c} \frac{p^2+1}{3} \\ \square \end{array} & \begin{array}{c} \frac{p^2+1}{3} \\ \square \end{array} & \begin{array}{c} \frac{2p^2}{3} \\ \square \end{array} & \begin{array}{c} \frac{p^2+1}{3} \\ \square \end{array} & \dots & \begin{array}{c} \frac{p^2+1}{3} \\ \square \end{array} \\ \hline \begin{array}{c} \frac{p^2+1}{3} \\ \square \end{array} & 0 & \frac{2}{3} & \frac{2}{3} & \frac{p^2+1}{3} & \frac{2}{3} & \dots & \frac{2}{3} \\ \begin{array}{c} \square \\ \frac{p^2+1}{3} \end{array} & \frac{2}{3} & 0 & \frac{2}{3} & \frac{p^2+1}{3} & \frac{2}{3} & \dots & \frac{p^2+1}{3} \\ \hline \begin{array}{c} \square \\ \frac{p^2+1}{3} \end{array} & \frac{2}{3} & \frac{2}{3} & 0 & \frac{p^2+1}{3} & \frac{2}{3} & \dots & \frac{2}{3} \\ \hline \begin{array}{c} \square \\ 2p^2 \end{array} & \begin{array}{c} p^2+1 \\ \square \end{array} & \begin{array}{c} p^2+1 \\ \square \end{array} & 0 & \begin{array}{c} \frac{p^2+1}{3} \\ \square \end{array} & \begin{array}{c} p^2+1 \\ \square \end{array} & \dots & \begin{array}{c} \frac{p^2+1}{3} \\ \square \end{array} \\ \hline \begin{array}{c} \square \\ \frac{p^2+1}{3} \end{array} & \frac{2}{3} & \frac{2}{3} & \frac{2}{3} & \frac{p^2+1}{3} & \frac{2}{3} & \dots & \frac{2}{3} \\ \hline \begin{array}{c} \square \\ \frac{p^2+1}{3} \end{array} & \begin{array}{c} p^2+1 \\ \square \end{array} & \begin{array}{c} \frac{2}{3} \\ \square \end{array} & \begin{array}{c} \frac{2}{3} \\ \square \end{array} & \begin{array}{c} \frac{2}{3} \\ \square \end{array} & \begin{array}{c} \frac{2}{3} \\ \square \end{array} & \dots & \begin{array}{c} \frac{2}{3} \\ \square \end{array} \end{array}.$$

$$\begin{array}{c} \text{h} \\ \psi - (-35p^2 - 129p + 162) \end{array} \begin{array}{c} \text{h} \\ \psi - (45p^2 - 103p + 122) \end{array} \begin{array}{c} \text{i} \\ [\psi - (-16p^2)] \end{array} \begin{array}{c} \text{i} \\ [\psi - (-16)] \end{array} (2p-3) = 0.$$

diam

$$\frac{2}{3} \quad \begin{matrix} \cdot \\ \cdot \\ \cdot \end{matrix} \quad \begin{matrix} \cdot \\ 0 \end{matrix}$$

[illegible]

–

Its characteristic polynomial is

*S
E
K
d
i
a
m*

$$\frac{(35p^2-129p+162)(45p^2-103p+122)(-16p^2)-16}{(2p-3)^2}.$$

$$\frac{1}{(2p-3)^2}.$$

$$\frac{E}{S} \frac{D}{d} i a m$$

$$-(35p^2-129p+162) \quad (45p^2-103p+122) \quad -16p^2$$

. 16 . . 16 . . 24 . .24 .

$$=\frac{1}{48}(272p^2-632p+756).$$

□

Theorem 4.5. Let F_p be a Friendship graph with p vertices, then

$$E_{diam}^{SSD}(F_p) = \sqrt{9760p^2 - 40544p + 43792 + 4(2p - 1)}.$$

Proof. The Friendship graph F_p with $2p+1$ vertices has its square sum degree by diameter matrix as follows

$$SSD_{diam}(F_p) = \begin{bmatrix} 0 & 2(p^2+1) & 2(p^2+1) & \dots & 2(p^2+1) \\ 2(p^2+1) & 0 & 4 & \dots & 4 \\ 2(p^2+1) & 4 & 0 & \ddots & 4 \\ 2(p^2+1) & 4 & \dots & 4 & 0 \end{bmatrix}.$$

Its characteristic polynomial is

$$\psi - (8p - 4) - \sqrt{9760p^2 - 40544p + 43792} \quad \psi - (8p - 4) - \sqrt{9760p^2 - 40544p + 43792} [\psi + (4)]^{2p-1} = 0$$

$$Spectra_{diam}^{SSD}(F_p) =$$

$$(8p - 4) + \sqrt{9760p^2 - 40544p + 43792} \quad (8p - 4) - \sqrt{9760p^2 - 40544p + 43792} \quad 4$$

$$\text{Therefore, } E_{diam}^{SSD}(F_p) =$$

$$\begin{aligned} & \cdot (8p - 4) + \sqrt{9760p^2 - 40544p + 43792} \cdot 1 + \cdot (8p - 4) - \sqrt{9760p^2 - 40544p + 43792} \cdot 1 + \cdot 4 \cdot (2p - 1) \\ & = \sqrt{9760p^2 - 40544p + 43792 + 4(2p - 1)}. \end{aligned}$$

□

References

- [1] B. Basavanagoud and E. Chitra. Degree square sum energy of graphs. *International Journal of Mathematics And its Applications* 6(2-B)(2018), 193205.
- [2] B. Basavanagoud and E. Chitra. Degree square sum polynomial of some special graphs. *International Journal of Applied Engineering Research*, 13(19)(2018), 1406014078.

- [3] B.BasavanagoudandE.Chitra.Degreeexponentsumenergyofagraph.
GulfJ.Math.,8(1)(2020),5270.
- [4]B. Basavanagoud and E. Chitra. Degree square sum equienergetic and hyperenergeticgraphs.*MalayaJ.Math.*, 8(2)(2020),301305.
- [5]I. Gutman. The energy of a graph.*Ber.Math.Stat. Sect. Forschungsz.Graz*.103(1978),122.
- [6]S. M. Hosamani and H. S. Ramane. On degree sum energy of a graph.
Eur.J.PureAppl.Math.9(3)(2016),340345.
- [7] G. Indulal,I.GutmanandA.Vijaykumar.Ondistanceenergyofgraphs.
MATCHCommun.Math.Comput.Chem..60(2)(2008),461472.
- [8]P.Mahalank,H.S.RamaneandA.R.Desai.Degreeexponentadjacencypolynomialofsomegraphs.*Adv.Math.,Sci*.J9(3)(2020),1001-1008.
- [9]H.S.Ramane,D.S.Revankar,I.Gutman,S.BhaskaraRao,B.D.AcharyaandH.B.Walikar.Boundsforthedistanceenergyofagraph.*KragujevacJ.Math*.31(2008),59-68
- [10] H.S.RamaneandS.S.Shinde.Degreeexponentpolynomialofgraphsobtained by some graph operations. *Electron. Notes Discrete Math*. 63(2017),161168.
- [11] H.S.RamaneandS.S.Shinde.Degreeexponentpolynomialanddegree exponent energy of graphs. *Indian Journal of Discrete Mathematics*2(1)(2016),17.
- [12] G. Polya and G. Szego, Problems and theorema in analysis, Series, Integralcalculus,Theoryoffunctions,*Springer*,(1972).
- [13] Gutman,I.(1978)TheEnergyofaGraph.*Ber.Math.Statist.Sekt.Forschungsz.Graz*,103,1-22.